

Adolescent Behaviors and Experiences Survey				
Sexual Behaviors	Female		Male	
	Percentage	Confidence Interval	Percentage	Confidence Interval
Ever had sexual intercourse	27.6	24.6 - 30.9	29.7	26.6 - 32.9
Had sexual intercourse for the first time before age 13 years	1.7	1.2 - 2.3	2.7	2.1 - 3.5
Had sexual intercourse with four or more persons during their life	4.8	3.7 - 6.2	6.8	5.4 - 8.5
Were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	19.5	17.2 - 22.1	18.0	15.5 - 20.9
Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	15.9	12.9 - 19.5	17.6	13.2 - 23.1
Used a condom during last sexual intercourse (among students who were currently sexually active)	48.3	43.4 - 53.1	60.9	56.0 - 65.5
Used birth control pills before last sexual intercourse with opposite-sex partner (to prevent pregnancy, not counting emergency contraception such as Plan B or the "morning after" pill, among students who were currently sexually active)	27.5	23.0 - 32.5	15.6	12.6 - 19.2
Were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)	6.5	5.1 - 8.3	5.1	4.3 - 6.1
Were tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	5.9	4.6 - 7.5	3.3	2.5 - 4.3
Used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	6.8	4.5 - 10.0	4.8	3.1 - 7.2
Used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	41.6	36.5 - 46.9	24.4	20.1 - 29.2
Used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	14.4	11.6 - 17.7	8.5	6.2 - 11.5
Did not use any method to prevent pregnancy during last sexual intercourse with an opposite-sex partner (among students who were currently sexually active)	14.1	10.8 - 18.2	12.6	9.2 - 16.9