

Table 26. Tuberculosis Cases and Percentages by Reason Tuberculosis Therapy Was Stopped and Type of Move: United States, 2014

Type of move	Total cases	Completed therapy		Adverse event		Lost		Refused		Died		Other ⁴		Unknown	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Moved in state ¹	289	240	(83.0)	1	(0.3)	4	(1.4)	3	(1.0)	15	(5.2)	24	(8.3)	2	(0.7)
Moved out of state ²	297	242	(81.5)	1	(0.3)	14	(4.7)	1	(0.3)	8	(2.7)	23	(7.7)	8	(2.7)
Moved out of country ³	373	165	(44.2)	1	(0.3)	19	(5.1)	3	(0.8)	11	(2.9)	166	(44.5)	7	(1.9)
Did not move	7,993	7,294	(91.3)	17	(0.2)	41	(0.5)	74	(0.9)	520	(6.5)	34	(0.4)	13	(0.2)
Unknown if moved	235	180	(76.6)	0	(0)	5	(2.1)	2	(0.9)	13	(5.5)	1	(0.4)	34	(14.5)

¹Includes patients who were alive at diagnosis, started on treatment, and moved in state.

²Includes patients who were alive at diagnosis, started on treatment, and moved out of state.

³Includes patients who were alive at diagnosis, started on treatment, and moved out of the country; transnational referrals were provided for 255 (68.4%) TB patients who moved out of the country.

⁴Therapy was discontinued for a known reason other than those listed (e.g., patient moved outside the United States, or patient moved from state A to state B, and although state A notified state B, state B never followed up).

Note: There may be differences in the way jurisdictions determine treatment completion for patients who moved out of the country; some reporting jurisdictions may be classifying all patients who moved out of the country as 'other' for reason therapy stopped.

Moving in and out of state or country is not mutually exclusive. Two patients moved twice, and one patient moved but did not indicate moving in state, out of state or out of country.